

**AMENDMENTS TO THE SPECIFICATION**

**Please replace the last paragraph on page 9 bridging page 10 with the following amended paragraph:**

As the photopolymerization initiator (B), for example, publicly known photopolymerization initiators can be used. Specific examples thereof include benzoin, benzoin methyl ether, benzoin ethyl ether, ~~benzyl benzil~~, diphenyl disulfide, tetramethylthiuram sulfide, diacetyl, eosin, thionine, Michlers' ketone, anthraquinone, chloroanthraquinone, methylantraquinone,  $\alpha$ -hydroxyisobutylphenone, p-isopropyl- $\alpha$ -hydroxyisobutylphenone,  $\alpha,\alpha'$ -dichloro-4-phenoxyacetophenone, 1-hydroxy-1-cyclohexylacetophenone, 2,2-dimethoxy-2-phenylacetophenone, methylbenzoyl formate, 2-methyl-1-[4-(methylthio)phenyl]-2-morpholinopropan-1-one, benzophenone, thioxanthone, 2-chlorothioxanthone, 2,4-diethylthioxanthone, 2-isopropylthioxanthone, 4-benzoyl-4'-methyldiphenyl sulfide, ethyl N,N-dimethylaminobenzoate, pentyl N,N-dimethylaminobenzoate, and triethanolamine. These photopolymerization initiators may be used alone, but are preferably used in combination.